

# Ponderosa Pine Forest Desired Conditions

Operational Draft: This document is prepared to provide guidance to Forest Plan revision teams. As this guidance is implemented, we expect to learn improved ways to do this work. As we learn, this document will be updated. This document was reviewed and revised as appropriate in April 2010 to conform to the requirements of the 1982 Planning Rule provisions.

## General Description

The ponderosa pine forest vegetation community includes two sub-types: Ponderosa pine bunchgrass and ponderosa pine Gambel oak. The ponderosa pine forest vegetation community generally occurs at elevations ranging from approximately 5,000 to 9,000 feet. It is dominated by ponderosa pine and commonly includes other species such as oak, juniper, and pinyon. More infrequently species such as aspen, Douglas-fir, white fir, and blue spruce may also be present, and may occur as individual trees. This forest vegetation community typically occurs with an understory of grasses and forbs although it sometimes includes shrubs.

## Landscape Scale Desired Conditions (10,000 + acres)

At the landscape scale, the ponderosa pine forest vegetation community is composed of trees from structural stages ranging from young to old. "Old growth" is well distributed in the landscape. Forest appearance is variable but generally uneven-aged and open; occasional areas of even-aged structure are present. The forest arrangement is in individual trees, small clumps, and groups of trees interspersed within variably-sized openings of grass/forbs/shrubs vegetation associations similar to historic patterns. Openings typically range from 10 percent in more productive sites to 70 percent in the less productive sites. Size, shape, number of trees per group, and number of groups per area are variable across the landscape. In the Gambel oak sub-type, all sizes and ages of oak trees are present. Denser tree conditions exist in some locations such as north facing slopes and canyon bottoms.

The ponderosa pine forest vegetation community is composed predominantly of vigorous trees, but declining trees are a component and provide for snags, top-killed, lightning- and fire-scarred trees, and coarse woody debris (>3 inch diameter), all well-distributed throughout the landscape. Ponderosa pine snags are typically 18 inches or greater at DBH and average 1 to 2 snags per acre. In the Gambel oak subtype, large oak snags (>10 inches) are a well-distributed component. Downed logs (>12 inch diameter at mid-point, >8 feet long) average 3 logs per acre within the forested area of the landscape. Coarse woody debris, including downed logs, ranges from 3 to 10 tons per acre.

The composition, structure, and function of vegetative conditions are resilient to the frequency, extent and severity of disturbances and climate variability. The landscape is a functioning ecosystem that contains all its components, processes, and conditions that result from endemic levels of disturbances (e.g. insects, diseases, fire, and wind), including snags, downed logs, and old trees. Grasses, forbs, shrubs, and needle cast (fine fuels), and small trees maintain the natural fire regime. Organic ground cover and herbaceous vegetation provide protection of soil, moisture infiltration, and contribute to plant and animal diversity and to ecosystem function. Frequent, low severity fires (Fire Regime I) are characteristic in this type, including throughout goshawk home ranges. Natural and anthropogenic disturbances are sufficient to maintain desired overall tree density, structure, species composition, coarse woody debris, and nutrient cycling.

## **Mid-Scale Desired Conditions (100 -1,000 acres)**

At the mid-scale the ponderosa pine forest vegetation community is characterized by variation in the size and number of tree groups depending on elevation, soil type, aspect, and site productivity. The more biologically productive sites contain more trees per group and more groups per area, resulting in less space between groups. Openings typically range from 10 percent in more productive sites to 70 percent in the less productive sites. Tree density within forested areas generally ranges from 20 to 80 square foot basal area per acre.

The mosaic of tree groups generally comprises an uneven-aged forest with all age classes present. Infrequently patches of even-aged forest structure are present. Disturbances sustain the overall age and structural distribution.

Fires burn primarily on the forest floor and do not spread between tree groups as crown fire.

Forest structure in the wildland urban interface (WUI)<sup>1</sup> can have smaller, more widely spaced groups of trees than in the non-WUI areas.

Forest conditions in goshawk post-fledging family areas (PFAs) are similar to general forest conditions except these forests contain 10 to 20 percent higher basal area in mid-aged to old tree groups than in goshawk foraging areas and the general forest. Goshawk nest areas have forest conditions that are multi-aged but are dominated by large trees with relatively dense canopies.

## **Fine Scale Desired Conditions ( ≤ 10acres)**

Trees typically occur in irregularly shaped groups and are variably-spaced with some tight clumps. Crowns of trees within the mid-aged to old groups are interlocking or nearly interlocking. Openings surrounding tree groups are variably-shaped and comprised of a grass/forb/shrub mix. Some openings contain individual trees. Trees within groups are of similar or variable ages and may contain species other than ponderosa pine. Size of tree groups typically is less than 1 acre, but averages .5 acres. Groups at the mid-aged to old stages consist of 2 to approximately 40 trees per group.

---

<sup>1</sup> Note –each Forest needs to provide the definition for WUI that they are using.